13. FOOTBALL

Priya B. Patel and Gerardo Miranda-Comas

■ HISTORY

Football emerged in the United States in the 1860s and 1870s as a direct consequence of the increasing importance of physical training and sport. Initially inspired by rugby in Great Britain, it soon rose to prominence and became America's "favorite pastime."¹

■ GOVERNING ORGANIZATIONS

- International: International Federation of American Football (IFAF)
- National: USA Football and National Football League (NFL)
- College: National Collegiate Athletic Association (NCAA)
- High School: National Federation of State High School Associations (NFHS)

■ PARTICIPANTS

Approximately 1.8 million athletes participate in American football in the United States.² These include athletes ranging from high school to professional level. In addition, there are more than 3 million participants in youth football leagues.³

■ RULES AND REGULATIONS

General rules are very similar throughout the different levels of play and leagues. At the collegiate and professional level, the total game time is 60 minutes split into four quarters of 15 minutes each and a break after the 2nd quarter (break time can vary). There are two 11-person teams that are in play. The offense has four downs to advance the ball at least 10 yards and if they are successful, they earn a first down, at which time they continue to try to advance another 10 yards toward the end zone or attempt a field goal. If they are unsuccessful, the ball is turned over to the other team to have a chance as the offense. Generally, a team scores by either making a touchdown or a field goal.

■ EQUIPMENT

The National Operating Committee on Standards for Athletic Equipment (NOCSAE) has specific requirements for the helmet. It should be fit one to two finger-breadths above eyebrows and the chin strap can be secured as a four-point or two-point system. The face mask provides easy visibility and others offer additional protection from facial injuries. As extra eye protection, some may also wear nontinted plexiglass shields. Tinted shields may be allowed with physician approval. Custom-made mouth guards are mandatory in college and high school. Shoulder pads vary with position, flat pads worn by those that need greater glenohumeral motion and cantilever pads for those who need more protection due to greater contact. Lower limb pads are used to protect against direct contact to the thighs.³

»»» **Adaptive Sport Key Points: Wheelchair Football**

Each team consists of six players and must include one female and one quadriplegic participant. The field measures 60 yards by 22 yards with 8 yards at each end zone. Games are split into two halves with each half consisting of 20 minutes. Halftime is 10 minutes. Play ends when the player holding the ball is tackled (a touch with one hand above the waist which includes the arm and hand).⁴

■ **MEDICAL COVERAGE LOGISTICS**

The role of the on-site physician leader is to organize and coordinate the medical staff, develop an emergency action plan (EAP), and interact with the coaches and officials to help ensure the safety of the players and spectators. Prior to the game, it is important to identify available medical staff, support personnel, field map, and the available medical equipment. Sports medicine physicians and Emergency Medical Services (EMS) personnel, along with sports medicine fellows, residents, medical students, athletic trainers, or physical therapists can attend expected medical needs. Establishing an EAP is the most important task. Protocols and guidelines should include role definition of the medical staff, emergency transportation including personnel, routes and adequate facilities with the capacity to evaluate and manage possible injuries, and an outline of the chain-of-command and communication standards. Contingency plans should be made for extreme temperature, humidity, and other weather conditions that may lead to game cancellation or postponement, or in game modifications like extra water breaks, especially at the youth level. Lastly, prior to the start of the competition, the medical team should confirm that players are wearing proper equipment. During the game the medical team and equipment is set up on the home team’s sideline. The referee will notify the head medical provider if an injured player needs to be evaluated on the field. Communication between individual members of the medical staff, coaches, referees, emergency medical staff, and local medical facilities is essential via two or more systems (e.g., hand-held radios and cell phones). Lastly, documentation is essential for review after the game/season to evaluate utilization and plan for future coverage.

■ **MEDICAL EMERGENCIES AND MEDICAL BAG ESSENTIALS**

Medical Bag Essentials in Football	
Medical Emergencies: Automated external defibrillator (AED) Tourniquet Cricothyrotomy kit 14- and 16-gauge needles Epi 1:1000 Diphenhydramine or another antihistamine Albuterol inhaler Rectal thermometer Ice bath Head and Spine: Concussion evaluation form Stretcher & Spine board Cervical orthosis Save a tooth kit Protective equipment removal tools	Fractures and Dislocations: Immobilizers and Slings Crutches Wound Care: Normal saline Betadine swabs Neosporin, mupirocin ointment Moleskin Elastic bandage Tube stretch gauze Sterile gauzes Vaseline Steri-strips Suture tray Lidocaine 1%

The collapsed or unconscious athlete is considered a medical emergency that requires activation of the proper channels of communication and protocols including basic life support (BLS)/advanced cardiac life support (ACLS). Although rarely reported, non-traumatic emergencies can include cardiac conditions like arrhythmias and hypertrophic cardiomyopathy (HCM), anaphylaxis, asthma, and heat-related illness, specifically heat stroke. Musculoskeletal injuries that may become medical emergencies include, upper and lower limb fractures or dislocations with neurovascular compromise, traumatic brain injuries with or without spinal cord involvement, and blunt thoracoabdominal trauma with possible internal bleeding.

■ EPIDEMIOLOGY

Football is a high velocity collision sport with a high incidence of traumatic and non-traumatic injuries. The rate of injuries in high school football players ranges from 3.2 to 8.1 per 1000 athlete exposures, and the rate of injuries in college players ranges from 3.1 to 11.1 per 1000 athlete exposures.² The lower limbs are the most commonly injured body area, specifically the knee followed by the ankle. Injuries to the shoulders and head/neck are also common.^{2,3}

■ COMMON INJURIES

Head and Spine Injuries

Head and spine injuries, especially **concussion**, are commonly seen in football at all levels. Mechanism of injury includes direct trauma to the head or due to an acceleration/deceleration force of the head due to trauma at another part of the body. Symptoms include confusion, dizziness, headaches, nausea and, in severe cases, loss of consciousness. Remove the player from play and conduct a neurological exam including a Sport Concussion Assessment Tool (SCAT) test.⁵ If no red flags are observed, then the player may remain on the sideline without his equipment. Periodic assessment should follow until cleared to leave with a parent if under age. He will need to follow up with a sports medicine provider in 24 to 48 hours for further assessment. Second impact syndrome is a possible complication if a concussed brain suffers another impact before resolution of the initial injury. **Cervical spine injury** was more commonly seen prior to new rules and regulations against tackling head first. Players who are found to be unconscious on the field or those reporting neck pain with neurologic symptoms from trauma during play should have the cervical spine immobilized. The recommended technique to transfer the injured player with a suspected cervical spine injury is the lift-and-slide technique which requires eight individuals: one at the head, three on each side and one controlling the spinal board. The other technique is the log roll and requires at least four providers: one controlling the head, two the body, and one the spinal board.⁶ Protective athletic equipment should be removed *prior* to transport to an emergency facility for an athlete-patient with suspected cervical spine instability. Equipment removal should be performed by at least three trained and experienced rescuers. If fewer than three people are present, the equipment should be removed at the earliest possible time after enough trained individuals arrive on the scene.⁶ They should be transported to the ED for further evaluation.

Upper Limb Injuries

Upper limb injuries include traumatic and overuse injuries. Common shoulder injuries include labral tears, bony injuries, and neurologic injuries. **Labral tears** commonly include a tear in the superior portion of the labrum and extend anterior to posterior also known as “SLAP” tears. Mechanism of injury is due to repetitive injuries to the

glenohumeral joint due to blocking or tackling. Symptoms include pain and weakness in the shoulder joint. Management includes removal from play; depending on the severity of injury, physical therapy and bracing may be used as well as surgery in some cases. **Bony injuries** include those to the glenoid labrum, known as a Bankart lesion, and to the humeral head, known as a Hills-Sachs lesion, often associated with glenohumeral joint (GHJ) dislocations and may require surgical management. **Brachial plexus neurapraxia**, also known as "stingers" or "burners," are due to a traction or compression of the brachial plexus. The mechanism of injury is usually the lateral flexion of the neck. Symptoms include unilateral burning pain, numbness or tingling down the arm, and weakness as well. These symptoms usually resolve quickly (minutes) and players can usually return to play after a thorough exam of the upper limb including range of motion and strength. Players with symptoms that do not resolve need further neurological testing to rule out cervical root avulsion. **Acromioclavicular (AC) joint** injuries are commonly seen in football and vary from sprain to tears of the ligaments around the AC joint. The mechanism of injury can be due to direct trauma to the lateral shoulder or falling on that point of the shoulder after a tackle. Symptoms include tenderness over the AC joint and you may feel a gap with displacement of the acromion from the clavicle. Players should be removed from play and obtain an x-ray of the shoulder. Depending on the severity of the injury, most will heal with conservative measures; however, some may require surgical intervention. **Mallet finger** is a flexion injury of the distal interphalangeal (DIP) joint. Mechanism of injury is due to trauma to the extensor tendon. Symptoms include difficulty to fully extend the DIP joint. Management includes obtaining x-rays to rule out a fracture and placing the DIP joint in mild hyperextension in splint for 6 weeks. **Jersey finger** is an avulsion injury of the flexor digitorum profundus. Mechanism of injury is due to extension of a flexed finger, usually when a player tries to grab a player for a tackle. Symptoms include inability to flex the DIP joint. Management includes surgical repair.

Lower Limb Injuries

The lower limbs are the most commonly injured body area in football. **Thigh contusion** is a common soft tissue injury in football. Mechanism of injury is usually blunt trauma. Symptoms include pain and swelling at the site. Ice and nonsteroidal anti-inflammatory medications (NSAIDs) are initially used. Players can return to play once strength is equal to the unaffected side. **Hip pointer** is a contusion of the iliac crest. Mechanism of injury is blunt trauma to the area. Symptoms include pain and inability to range the hip. Patient should be removed from play and immediately start icing area of pain. **Medial collateral ligament (MCL)** injury is the most common knee injury in football. Mechanism of action is a valgus load to the knee by direct trauma usually through tackling. Symptoms vary by grades of injury (Grades I–III), with Grade II as a partial ligament injury and Grade III as a full ligament tear. Valgus laxity testing at the knee will reveal if there is laxity with or without an end point. Management includes conservative treatment (rest, ice, NSAIDs). Some players may wear protective bracing. MCL injuries may occur concomitantly with **Anterior cruciate ligament (ACL)** and/or **medial meniscus tears** as part of the "unhappy triad." ACL tears can occur secondary to a direct valgus load, or without contact during a single leg landing from a jump, or during a cutting move while the foot is planted. An acute knee injury with immediate swelling after one of the mechanisms of injury mentioned, correlates with an ACL tear. The player cannot continue playing during the game and will need further imaging, MRI is the gold standard and possible surgical intervention. **Hamstring strain** is commonly seen due to noncontact sudden muscle contraction. Symptoms include tenderness over the area and players may have edema and ecchymosis as well. Management

includes rest and return to play when patient is asymptomatic. **Lisfranc fracture** is commonly seen as a running injury. Mechanism of injury is due to twisting of the foot when a player is falling. Symptoms include pain and inability to ambulate on the affected foot. Player should be removed from play, sent to ED to obtain an x-ray. Management includes open reduction internal fixation (ORIF) surgery. **Turf toe** is an injury to the first metatarsophalangeal (MTP) joint. Mechanism of injury is due to trauma causing dorsiflexion of the planted toe. Symptoms include pain and swelling at the area and painful ambulation as a result. Management includes rigid shoe insert, donut padding, rest, and ice. Players can return to play when they are pain free.

■ SUMMARY

Football is a popular high-velocity contact sport that can be played at all levels. Understanding the game, positions, and most common injuries is essential for sideline coverage.

■ References

1. Collins T. Unexceptional exceptionalism: The origins of American football in a transnational context. *J Glob Hist.* 2013;8(2):209-230. doi:10.1017/S1740022813000193.
2. Shah S, Luftman JP, Vigil DV. Football: Sideline management of injuries. *Curr Sports Med Rep.* 2004;3(3):146-153. doi:10.1007/s11932-004-0062-5.
3. Rizzone K, Diamond A, Gregory A. Sideline coverage of youth football. *Curr Sports Med Rep.* 2013;12(3):143-149. doi:10.1249/JSR.0b013e3182955d1c.
4. Wheelchair Sports Federation. Football. Updated December 11, 2009. <http://wheelchairsportsfederation.org/adaptive-sports/football>
5. McCrory P, Meeuwisse W, Dvorak J, et al. Consensus statement on concussion in sport—the 5th International Conference on Concussion in Sport held in Berlin, October 2016. *Br J Sports Med.* 2017;13(2):53-65. doi:10.1136/bjsports-2017-097699.
6. National Athletic Trainers' Association. Appropriate care of the spine injured athlete. Updated from 1998 document. Prof Rep. 2015. <https://www.nata.org/blog/toddc/update-appropriate-care-spine-injured-athlete>